α					4	\sim
·.	$^{\circ}$ 1	er	0	Δ		<i>1</i> 1
. 7			ı			ι,

Name:		
Block:	Date:	

Walk This Way

Purpose:

To determine the average velocity of several individuals moving at different speeds and directions.

Procedure:

See Pages 372-3 in "BC Science 10" (Conduct an Investigation 8-2E "Walk This Way") Note: additional trials will be recorded as indicated by the data tables below.

Data Tables:

Walking Forward											
Position (m)	0	5	10	15	20	25	30	35	40	45	50
Time (s)	0	3.06	6.52	רניסו	12.61	15.7	18.9	21.96	25.06	28.59	31.5

Jogging Forward											
Position (m)	0	5	10	15	20	25	30	35	40	45	50
Time (s)	0	2.0	3.65	5.34	6.71	8.36	9.9	11.58	12.87	14.75	16.8

Running Forward											
Position (m)	0	5	10	15	20	25	30	35	40	45	50
Time (s)	0	1.33	2.03	2.25	3,41	4.01	4.6	5,33	5.17	6.67	7.5

Walking Backward											
Position (m)	50	45	40	35	30	25	20	15	10	5	0
Time (s)	0	3.51	6.62	10.19	1371	17.01	21.66	25.36	28,61	3273	35.8

~	•	10
V. C	ience	
L) L		, 11

Name:	
Block:	Date:

Analysis of Results:

- 1. On the same set of axes, plot a position-time graph for all 4 sets of data.
- 2. On the graph:
 - a. Draw a best-fit line for each set of data.
 - b. Calculate the slope of each of the best fit lines. (make sure to include units)
- 3. What is the average velocity of the student...
 - a. Walking forward?
 - b. Jogging forward?
 - c. Running forward?
 - d. Walking backward?
- 4. Was the average speed of the student faster when walking forward or backward?
- 5. Did any of the students have perfectly uniform motion while walking, jogging or running forward or backward? Use your graph to justify your answer.

Conclusion:

What can you infer about the slope of a position-time graph of an object's and its average velocity? Include comments about the steepness of the slope and whether it is positive or negative.